

Lecture 15

Tues. 3.08.2016

Introduction to classical oscillation and resonance

(Ch. 2 of Unit 2)

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

*Linear **damped**-harmonic oscillator equation of motion.*

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

*Linear **forced-damped**-harmonic oscillator equation of motion.*

*Properties of **Green's function** solutions and their mathematical/physical behavior*

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

*Complete **Green's Solution** for the **FDHO** (**Forced-Damped-Harmonic Oscillator**)*

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

*Approximate Lorentz-**Green's Function** for high quality **FDHO** (Quantum propagator)*

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

➔ *1D forced-damped-harmonic oscillator equations and Green's function solutions*

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)


Smith Charts

Linear forced-damped-harmonic oscillator equation of motion.

$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$

Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)



$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$

Coordinate $z=z(t)$ is the response coordinate for a particle of mass m and charge e

driven by external **stimulating force** $\longrightarrow F_{stimulus}(t) = eE(t)$

held back by a **harmonic (linear) restoring force** $\longrightarrow F_{restore} = -kz, (k = \omega_0^2 m),$

retarded by **frictional damping force** $\longrightarrow F_{damping} = -b \frac{dz}{dt}, (b = 2\Gamma m)$

1D forced-damped-harmonic oscillator equations and Green's function solutions



Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)


Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Linear

harmonic oscillator equation of motion.

Phasor Clock
 $Z(t) = Ae^{-i(\omega t)}$
 $= A\cos(\omega t)$
 $-i A\sin(\omega t)$
 Phasor clocks
 turn
 clockwise
 in time for
 positive ω



$$F_{total}(t) = m \frac{d^2 z}{dt^2} =$$

$F_{restore}$

$$\frac{d^2 z}{dt^2} =$$

$$\frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + \omega_0^2 z = 0$$

$$+ \omega_0^2 z = 0$$


Coordinate $z=z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a harmonic (linear) restoring force

$$F_{restore} = -kz, \quad (k = \omega_0^2 m),$$

Linear harmonic oscillator equation of motion.

Fig. 2.2.1




$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + \omega_0^2 z = 0$$

Coordinate $z=z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a harmonic (linear) restoring force $\longrightarrow F_{restore} = -kz, (k = \omega_0^2 m),$



The famous one-Hertz ($\nu_0=1/s.$ or: $\omega_0 = 2\pi = 6.2832rad/s.$) oscillator.

[OscillIt Web Simulation: Phasor description of Harmonic Oscillation](#)

Fig. 2.2.2 Phasor z and corresponding coordinate versus time plot for $\omega_0=2\pi$ and $\Gamma=0$

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

➔ *Linear damped-harmonic oscillator equation of motion.*

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty


Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Linear *damped-harmonic oscillator equation of motion.*

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$


Coordinate $z=z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a **harmonic (linear) restoring force** $\longrightarrow F_{restore} = -kz, (k = \omega_0^2 m),$
 retarded by **frictional damping force** $\longrightarrow F_{damping} = -b \frac{dz}{dt}, (b = 2\Gamma m)$

[OscillIt Web Simulation: Phasor description of Harmonic Oscillation](#)

Linear damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$

Trick:
Set: $z = z(t) = Ae^{-i\omega t}$

$$\begin{aligned} [(-i\omega)^2 + 2\Gamma(-i\omega) + \omega_0^2] e^{-i\omega t} &= 0 \\ \omega^2 + 2i\Gamma\omega - \omega_0^2 &= 0 \end{aligned}$$

Coordinate $z = z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a harmonic (linear) restoring force

$$F_{restore} = -kz$$


retarded by frictional damping force

$$F_{damping} = -b \frac{dz}{dt}$$

[OscillIt Web Simulation: Phasor description of Harmonic Oscillation](#)

Linear damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$

Trick:
 Set: $z = z(t) = Ae^{-i\omega t}$

$$\left[(-i\omega)^2 + 2\Gamma(-i\omega) + \omega_0^2 \right] e^{-i\omega t} = 0$$

$$\omega^2 + 2i\Gamma\omega - \omega_0^2 = 0$$

Solve for: $\omega = \omega_{\pm}$

$$\omega_{\pm} = \frac{-2i\Gamma \pm \sqrt{-4\Gamma^2 + 4\omega_0^2}}{2}$$

Coordinate $z = z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a harmonic (linear) restoring force

$$F_{restore} = -kz$$


retarded by frictional damping force

$$F_{damping} = -b \frac{dz}{dt}$$

[OscillIt Web Simulation: Phasor description of Harmonic Oscillation](#)

Linear damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$

Trick:
Set: $z = z(t) = Ae^{-i\omega t}$

$$[(-i\omega)^2 + 2\Gamma(-i\omega) + \omega_0^2] e^{-i\omega t} = 0$$

$$\omega^2 + 2i\Gamma\omega - \omega_0^2 = 0$$

Solve for: $\omega = \omega_{\pm}$

$$\omega_{\pm} = \frac{-2i\Gamma \pm \sqrt{-4\Gamma^2 + 4\omega_0^2}}{2}$$

$$= -i\Gamma \pm \sqrt{\omega_0^2 - \Gamma^2}$$

Coordinate $z = z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a harmonic (linear) restoring force \longrightarrow


$$F_{restore} = -kz$$

retarded by frictional damping force \longrightarrow

$$F_{damping} = -b \frac{dz}{dt}$$

Linear damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$

Trick:
Set: $z = z(t) = Ae^{-i\omega t}$

$$\left[(-i\omega)^2 + 2\Gamma(-i\omega) + \omega_0^2 \right] e^{-i\omega t} = 0$$

$$\omega^2 + 2i\Gamma\omega - \omega_0^2 = 0$$

Solve for: $\omega = \omega_{\pm}$

$$\omega_{\pm} = \frac{-2i\Gamma \pm \sqrt{-4\Gamma^2 + 4\omega_0^2}}{2}$$

$$= -i\Gamma \pm \sqrt{\omega_0^2 - \Gamma^2}$$

Solution:

$$z(t) = e^{-i\left(-i\Gamma \pm \sqrt{\omega_0^2 - \Gamma^2}\right)t}$$

$$= e^{\left(-\Gamma \pm i\sqrt{\omega_0^2 - \Gamma^2}\right)t}$$

$$= e^{-\Gamma t} e^{\pm i\sqrt{\omega_0^2 - \Gamma^2}t}$$

$$= e^{-\Gamma t} e^{\pm i\omega_{\Gamma}t}$$

Coordinate $z = z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a harmonic (linear) restoring force \longrightarrow

$$F_{restore} = -kz$$

retarded by frictional damping force \longrightarrow

$$F_{damping} = -b \frac{dz}{dt}$$

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.



Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty


Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Linear *damped-harmonic oscillator equation of motion.*

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$

Trick:
Set: $z = z(t) = A e^{-i\omega t}$

$$\left[(-i\omega)^2 + 2\Gamma(-i\omega) + \omega_0^2 \right] e^{-i\omega t} = 0$$

$$\omega^2 + 2i\Gamma\omega - \omega_0^2 = 0$$

Solve for: $\omega = \omega_{\pm}$

$$\omega_{\pm} = \frac{-2i\Gamma \pm \sqrt{-4\Gamma^2 + 4\omega_0^2}}{2}$$

$$= -i\Gamma \pm \sqrt{\omega_0^2 - \Gamma^2}$$

Solution:

$$z(t) = e^{-i \left(-i\Gamma \pm \sqrt{\omega_0^2 - \Gamma^2} \right) t}$$

$$= e^{\left(-\Gamma \pm i\sqrt{\omega_0^2 - \Gamma^2} \right) t}$$

$$= e^{-\Gamma t} e^{\pm i\sqrt{\omega_0^2 - \Gamma^2} t}$$

$$= e^{-\Gamma t} e^{\pm i\omega_{\Gamma} t}$$

Coordinate $z = z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a **harmonic (linear) restoring force** \longrightarrow

$$F_{restore} = -kz$$

retarded by **frictional damping force** \longrightarrow


$$F_{damping} = -b \frac{dz}{dt}$$

It oscillates at an angular frequency ω_{Γ} reduced slightly by .05% from ω_0 due to damping $\Gamma = 0.2$.

$$\omega_{\Gamma} = \sqrt{\omega_0^2 - \Gamma^2} = \omega_0 - \frac{1}{2} (\Gamma^2 / \omega_0) + \dots = 6.2831853 - 0.003183 + \dots = 6.280002 + \dots = 6.280001$$

Linear *damped-harmonic oscillator equation of motion.*

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$

Trick:
Set: $z = z(t) = Ae^{-i\omega t}$

$$\left[(-i\omega)^2 + 2\Gamma(-i\omega) + \omega_0^2 \right] e^{-i\omega t} = 0$$

$$\omega^2 + 2i\Gamma\omega - \omega_0^2 = 0$$

Solve for: $\omega = \omega_{\pm}$

$$\omega_{\pm} = \frac{-2i\Gamma \pm \sqrt{-4\Gamma^2 + 4\omega_0^2}}{2}$$

$$= -i\Gamma \pm \sqrt{\omega_0^2 - \Gamma^2}$$

Solution:

$$z(t) = e^{-i \left(-i\Gamma \pm \sqrt{\omega_0^2 - \Gamma^2} \right) t}$$

$$= e^{\left(-\Gamma \pm i\sqrt{\omega_0^2 - \Gamma^2} \right) t}$$

$$= e^{-\Gamma t} e^{\pm i\sqrt{\omega_0^2 - \Gamma^2} t}$$

$$= e^{-\Gamma t} e^{\pm i\omega_{\Gamma} t}$$


Coordinate $z = z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a harmonic (linear) restoring force

$$F_{restore} = -kz$$

retarded by frictional damping force

$$F_{damping} = -b \frac{dz}{dt}$$



[OscillIt Web Simulation: Phasor description of Damped Harmonic Oscillation](#)

Fig. 2.2.3 Phasor z and corresponding coordinate versus time plot for $\omega_0 = 2\pi$ and $\Gamma = 0.2$

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

➔ *Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)*

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty


Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Linear *damped-harmonic oscillator equation of motion.*

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$

Oscillator Figures of Merit:

Number N of oscillations to reduce amplitude to 5%

Easy-to-recall 5% approximation:

$$e^{-3} \cong 0.05$$


Coordinate $z=z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a **harmonic (linear) restoring force**

$$F_{restore} = -kz$$

retarded by **frictional damping force**

$$F_{damping} = -b \frac{dz}{dt}$$




$$t_{5\%} = \frac{3}{\Gamma} = \frac{3}{0.2} = 15$$

[OscillIt Web Simulation: Phasor description of Damped Harmonic Oscillation](#)

Fig. 2.2.3 Phasor z and corresponding coordinate versus time plot for $\omega_0=2\pi$ and $\Gamma=0.2$

Linear *damped-harmonic oscillator equation of motion.*

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m}$$

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = 0$$

Oscillator Figures of Merit:

Number N of oscillations to reduce amplitude to 5% (or 4.321%)

Easy-to-recall 5% approximation:

$$e^{-3} \cong 0.05$$

More precise one:

$$e^{-\pi} \cong 0.04321$$

$$N_{5\%} = \frac{\omega_\Gamma \cdot t_{5\%}}{2\pi} = \frac{3\omega_\Gamma}{2\pi\Gamma} \sim \frac{\omega_\Gamma}{2\Gamma}$$

$$t_{4.321\%} = \frac{\pi}{\Gamma} = \frac{\pi}{0.2} = 15.708$$

Coordinate $z=z(t)$ is the response coordinate for a particle of mass m and charge e

held back by a harmonic (linear) restoring force $\longrightarrow F_{restore} = -kz$

retarded by frictional damping force $\longrightarrow F_{damping} = -b \frac{dz}{dt}$




Fig. 2.2.3 Phasor z and corresponding coordinate versus time plot for $\omega_0=2\pi$ and $\Gamma=0.2$

[OscillIt Web Simulation: Phasor description of Damped Harmonic Oscillation](#)

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

➔ *Linear forced-damped-harmonic oscillator equation of motion.*

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty


Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Linear forced-damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$

Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$

Coordinate $z=z(t)$ is the response coordinate for a particle of mass m and charge e


driven by external **stimulating force** $\longrightarrow F_{stimulus}(t) = eE(t)$

held back by a **harmonic (linear) restoring force** $\longrightarrow F_{restore} = -kz, (k = \omega_0^2 m),$

retarded by **frictional damping force** $\longrightarrow F_{damping} = -b \frac{dz}{dt}, (b = 2\Gamma m)$

Linear forced-damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$


Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$

Solving for $z_{stimulus}(t)$ given $a_{stimulus}$:

Linear forced-damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$

Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$


Solving for $z_{stimulus}(t)$ given $a_{stimulus}$:

$$\left(\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2 \right) z = a_{stimulus}$$

$$z = \frac{1}{\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2} a_{stimulus}$$

Linear forced-damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$

Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$

$$\left(\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2 \right) z = a_{stimulus}$$


$$z = \frac{1}{\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2} a_{stimulus}$$

Solving for $z_{stimulus}(t)$ given $a_{stimulus}$:

Pretty crazy?

Linear forced-damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$

Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$

Solving for $z_{stimulus}(t)$ given $a_{stimulus}$:

$$\left(\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2 \right) z = a_{stimulus}$$


Pretty crazy? But not so crazy if

$$a_{stimulus}(t) = |a_{stimulus}| e^{-i\omega_{stimulus} t} = |a_s| e^{-i\omega_s t}$$

$$z = \frac{1}{\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2} a_{stimulus}$$

Linear forced-damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$

Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$

$$\left(\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2 \right) z = a_{stimulus}$$

Solving for $z_{stimulus}(t)$ given $a_{stimulus}$:


Pretty crazy? But not so crazy if $a_{stimulus}(t) = |a_{stimulus}| e^{-i\omega_{stimulus} t} = |a_s| e^{-i\omega_s t}$

$$z_{stimulus} = \frac{1}{-\omega_s^2 - i2\Gamma\omega_s + \omega_0^2} a_s e^{-i\omega_s t}$$

$$z_s e^{-i\omega_s t} = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} a_s e^{-i\omega_s t}$$

Linear forced-damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$

Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$

Solving for $z_{stimulus}(t)$ given $a_{stimulus}$:

$$\left(\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2 \right) z = a_{stimulus}$$

Pretty crazy? But not so crazy if

$$a_{stimulus}(t) = |a_{stimulus}| e^{-i\omega_{stimulus} t} = |a_s| e^{-i\omega_s t}$$


$$z_{stimulus} = \frac{1}{-\omega_s^2 - i2\Gamma\omega_s + \omega_0^2} a_s e^{-i\omega_s t}$$

$$z_s e^{-i\omega_s t} = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} a_s e^{-i\omega_s t}$$

$$z_s = G_{\omega_0}(\omega_s) \cdot a_s$$

Linear forced-damped-harmonic oscillator equation of motion.

Fig. 2.2.1



$$F_{total}(t) = m \frac{d^2 z}{dt^2} = F_{damping} + F_{restore} + F_{stimulus}$$

$$\frac{d^2 z}{dt^2} = \frac{F_{damping}}{m} + \frac{F_{restore}}{m} + \frac{F_{stimulus}}{m}$$

Stimulating acceleration $a_{stimulus} = a(t)$ due to stimulating force $F_{stimulus}(t)$ (Typically **E**-field)

$$\frac{d^2 z}{dt^2} + 2\Gamma \frac{dz}{dt} + \omega_0^2 z = a_{stimulus} = \frac{e}{m} E(t)$$

$$\left(\frac{d^2}{dt^2} + 2\Gamma \frac{d}{dt} + \omega_0^2 \right) z = a_{stimulus}$$

Solving for $z_{stimulus}(t)$ given $a_{stimulus}$:

$$z_{stimulus} = \frac{1}{-\omega_s^2 - i2\Gamma\omega_s + \omega_0^2} a_s e^{-i\omega_s t}$$

Pretty crazy? But not so crazy if $a_{stimulus}(t) = |a_{stimulus}| e^{-i\omega_{stimulus} t} = |a_s| e^{-i\omega_s t}$

$$z_s e^{-i\omega_s t} = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} a_s e^{-i\omega_s t}$$

$$z_s = G_{\omega_0}(\omega_s) \cdot a_s$$

Green's Function for the F-D-H Oscillator (FDHO)

George Green (14 July 1793 – 31 May 1841)

[Mathematical Analysis to the Theories of Electricity and Magnetism \(1828\)](#)

[Wiki](#)

[The Green of Green's Function - Physics Today \(2003 Dec\)](#)

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

 *Properties of Green's function solutions and their mathematical/physical behavior*

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Green's Function for the FDHO (Forced-Damped-Harmonic Oscillator)




Fig. 2.2.4 Black-box diagram of oscillator response to monochromatic stimulus

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} = \text{Re } G_{\omega_0}(\omega_s) + i \text{Im } G_{\omega_0}(\omega_s)$$

Real and imaginary parts of the *rectangular form* of G :

Green's Function for the FDHO (Forced-Damped-Harmonic Oscillator)




Fig. 2.2.4 Black-box diagram of oscillator response to monochromatic stimulus

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} = \text{Re } G_{\omega_0}(\omega_s) + i \text{Im } G_{\omega_0}(\omega_s)$$

Real and imaginary parts of the *rectangular form* of G : $\frac{1}{x-iy} = \frac{1}{x-iy} \frac{x+iy}{x+iy} = \frac{x+iy}{x^2+y^2}$

Green's Function for the FDHO (Forced-Damped-Harmonic Oscillator)




Fig. 2.2.4 Black-box diagram of oscillator response to monochromatic stimulus

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} = \text{Re } G_{\omega_0}(\omega_s) + i \text{Im } G_{\omega_0}(\omega_s)$$

Real and imaginary parts of the *rectangular form* of G : $\frac{1}{x-iy} = \frac{1}{x-iy} \frac{x+iy}{x+iy} = \frac{x+iy}{x^2+y^2} = \frac{x}{x^2+y^2} + i \frac{y}{x^2+y^2}$

$$\text{Re } G_{\omega_0}(\omega_s) = \frac{\omega_0^2 - \omega_s^2}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

$$\text{Im } G_{\omega_0}(\omega_s) = \frac{2\Gamma\omega_s}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

Green's Function for the FDHO (Forced-Damped-Harmonic Oscillator)




Fig. 2.2.4 Black-box diagram of oscillator response to monochromatic stimulus

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} = \text{Re } G_{\omega_0}(\omega_s) + i \text{Im } G_{\omega_0}(\omega_s) = |G_{\omega_0}(\omega_s)| e^{i\rho}$$

Real and imaginary parts of the *rectangular form* of G :

$$\text{Re } G_{\omega_0}(\omega_s) = \frac{\omega_0^2 - \omega_s^2}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

$$\text{Im } G_{\omega_0}(\omega_s) = \frac{2\Gamma\omega_s}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

Magnitude $|G_{\omega_0}(\omega_s)|$ and polar angle ρ of the *polar form* of G :

$$|G_{\omega_0}(\omega_s)| = \frac{1}{\sqrt{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}}$$

$$\rho = \tan^{-1} \left(\frac{2\Gamma\omega_s}{\omega_0^2 - \omega_s^2} \right)$$

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior



Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Green's Function for the FDHO (Forced-Damped-Harmonic Oscillator)




Fig. 2.2.4 Black-box diagram of oscillator response to monochromatic stimulus

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} = \text{Re } G_{\omega_0}(\omega_s) + i \text{Im } G_{\omega_0}(\omega_s) = |G_{\omega_0}(\omega_s)| e^{i\rho}$$

Real and imaginary parts of the *rectangular form* of G :

$$\text{Re } G_{\omega_0}(\omega_s) = \frac{\omega_0^2 - \omega_s^2}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$



$$\text{Im } G_{\omega_0}(\omega_s) = \frac{2\Gamma\omega_s}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

Magnitude $|G_{\omega_0}(\omega_s)|$ and *polar angle* ρ of the *polar form* of G :

$$|G_{\omega_0}(\omega_s)| = \frac{1}{\sqrt{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}}$$

$$\rho = \tan^{-1}\left(\frac{2\Gamma\omega_s}{\omega_0^2 - \omega_s^2}\right)$$

polar angle ρ is the *phase lag angle* ρ



$$z_{\text{response}}(t) = |G_{\omega_0}(\omega_s)| a(0) e^{-i(\omega_s t - \rho)}$$

Fig. 2.2.5 Oscillator response and stimulus phasors rotate rigidly at angular rate ω_s .

Green's Function for the FDHO (Forced-Damped-Harmonic Oscillator)




Fig. 2.2.4 Black-box diagram of oscillator response to monochromatic stimulus

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} = \text{Re } G_{\omega_0}(\omega_s) + i \text{Im } G_{\omega_0}(\omega_s) = |G_{\omega_0}(\omega_s)| e^{i\rho}$$

Real and imaginary parts of the *rectangular form* of G :

$$\text{Re } G_{\omega_0}(\omega_s) = \frac{\omega_0^2 - \omega_s^2}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

$$\text{Im } G_{\omega_0}(\omega_s) = \frac{2\Gamma\omega_s}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

Magnitude $|G_{\omega_0}(\omega_s)|$ and *polar angle* ρ of the *polar form* of G :

$$|G_{\omega_0}(\omega_s)| = \frac{1}{\sqrt{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}}$$

$$\rho = \tan^{-1}\left(\frac{2\Gamma\omega_s}{\omega_0^2 - \omega_s^2}\right)$$

polar angle ρ is the *phase lag angle* ρ





Fig. 2.2.5 Oscillator response and stimulus phasors rotate rigidly at angular rate ω_s .

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification



Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)


Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts



OscillIt Web Simulation
 Lorentz Response

Fig. 2.2.6 Anatomy of oscillator Green-Lorentz response function plots


Phase lag angle

$$\rho = \tan^{-1} \left(\frac{2\Gamma\omega_s}{\omega_0^2 - \omega_s^2} \right)$$

$$\text{Re } G_{\omega_0}(\omega_s) = \frac{\omega_0^2 - \omega_s^2}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

$$\text{Im } G_{\omega_0}(\omega_s) = \frac{2\Gamma\omega_s}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

$$AAF = \frac{\text{Resonant response}}{\text{DC response}} = \frac{|G_{\omega_0}(\omega_s = \omega_0)|}{|G_{\omega_0}(0)|} = \frac{1/(2\Gamma\omega_0)}{1/\omega_0^2} = \frac{\omega_0}{2\Gamma} \equiv q \quad (\text{angular quality factor})$$



[OscillIt Web Simulation](http://www.uark.edu/ua/modphys/markup/OscillItWeb.html)
 Lorentz Response ($\Gamma=0.2$)

Fig. 2.2.6 Anatomy of oscillator Green-Lorentz response function plots

Phase lag angle

$$\rho = \tan^{-1} \left(\frac{2\Gamma\omega_s}{\omega_0^2 - \omega_s^2} \right)$$

$$\text{Re } G_{\omega_0}(\omega_s) = \frac{\omega_0^2 - \omega_s^2}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

Real part

$$\text{Im } G_{\omega_0}(\omega_s) = \frac{2\Gamma\omega_s}{(\omega_0^2 - \omega_s^2)^2 + (2\Gamma\omega_s)^2}$$

Imaginary part

$$AAF = \frac{\text{Resonant response}}{\text{DC response}} = \frac{|G_{\omega_0}(\omega_s = \omega_0)|}{|G_{\omega_0}(0)|} = \frac{1/(2\Gamma\omega_0)}{1/\omega_0^2} = \frac{\omega_0}{2\Gamma} \equiv q \quad (\text{angular quality factor})$$

OscillIt Web Simulation
Lorentz Response ($\Gamma=0.2$)

OscillIt Web Simulation
Lorentz Response ($\Gamma=0.1$)




Fig. 2.2.7 Comparing Lorentz-Green resonance region for (a) $\Gamma=0.2$ and (b) $\Gamma=0.1$.

Maximum and minimum points of $\text{Re}G(\omega)$ and inflection points of $\text{Im}G(\omega)$ are near region boundaries $\omega^{\text{FWHM}(\pm)} = \omega_0 \pm \Gamma$.

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

➔ *Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)*

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Complete *Green's Solution* for the *FDHO* (*Forced-Damped-Harmonic Oscillator*)

$$\begin{aligned} z(t) &= z_{\text{transient}}(t) + z_{\text{response}}(t) \equiv z_{\text{decaying}}(t) + z_{\text{steady state}}(t) \\ &= Ae^{-\Gamma t} e^{-i\omega_{\Gamma} t} + G_{\omega_0}(\omega_s) a(0) e^{-i\omega_s t} \\ &= Ae^{-\Gamma t} e^{-i\omega_{\Gamma} t} + \left| G_{\omega_0}(\omega_s) \right| a(0) e^{-i(\omega_s t - \rho)} \end{aligned}$$

Known as “homogeneous” solution (no force)
Let's you set initial values or boundary conditions

Known as “inhomogeneous” solution
Not function of initial values. Marches to stimulus only.

Complete *Green's Solution* for the *FDHO* (*Forced-Damped-Harmonic Oscillator*)

$$\begin{aligned}z(t) &= z_{\text{transient}}(t) + z_{\text{response}}(t) \equiv z_{\text{decaying}}(t) + z_{\text{steady state}}(t) \\ &= Ae^{-\Gamma t} e^{-i\omega_{\Gamma} t} + G_{\omega_0}(\omega_s) a(0) e^{-i\omega_s t} \\ &= Ae^{-\Gamma t} e^{-i\omega_{\Gamma} t} + \left| G_{\omega_0}(\omega_s) \right| a(0) e^{-i(\omega_s t - \rho)}\end{aligned}$$

Known as “homogeneous” solution (no force)
Let's you set initial values or boundary conditions

Known as *Transient* solution since it dies-off as time
advances past initial conditions

Known as “inhomogeneous” solution
Not function of initial values. Marches to stimulus only.

Known as *Steady State* solution since it is present as long as stimulus is.

Complete *Green's Solution* for the *FDHO* (*Forced-Damped-Harmonic Oscillator*)

$$\begin{aligned}
 z(t) &= z_{\text{transient}}(t) + z_{\text{response}}(t) \equiv z_{\text{decaying}}(t) + z_{\text{steady state}}(t) \\
 &= Ae^{-\Gamma t} e^{-i\omega_{\Gamma} t} + G_{\omega_0}(\omega_s) a(0) e^{-i\omega_s t} \\
 &= Ae^{-\Gamma t} e^{-i\omega_{\Gamma} t} + \left| G_{\omega_0}(\omega_s) \right| a(0) e^{-i(\omega_s t - \rho)}
 \end{aligned}$$

Known as “homogeneous” solution (no force)
 Let's you set initial values or boundary conditions

Known as *Transient* solution since it dies-off as time advances past initial conditions


Known as “inhomogeneous” solution
 Not function of initial values. Marches to stimulus only.

Known as *Steady State* solution since it is present as long as stimulus is.

Stimulus: $A_s = 0.5000$ $\omega = 6.2832$
 Response: $R = 0.1989$ $\rho = 1.5708$

About $t = 3/\Gamma = 15 \text{ sec}$

About $t = \text{forever}$



OscillIt Web Simulation FDHO - Driven on resonance

Fig. 2.2.8 On Resonance (a) Response z -phasor lags $\rho = 90^\circ$ behind stimulus F -phasor. ($\omega_s = \omega_0 = 2\pi$, $\omega_0 = 2\pi$, and $\Gamma = 0.2$). (b) Time plots of $\text{Re } z(t)$ and $\text{Re } F(t)$

Fig. 2.2.8 Below Resonance (c) Response z -phasor lags $\rho = 8.05^\circ$ behind stimulus F -phasor. ($\omega_s = 5.03$, $\omega_0 = 2\pi$, and $\Gamma = 0.2$). (d) Time plots of $\text{Re } z(t)$ and $\text{Re } F(t)$. Beats are barely visible.

OscillIt Web Simulation FDHO - Driven well below resonance

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)


Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)



Transient solutions vs. Steady State solutions


Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts


Stimulus: $A_s = 1.0000$ $\omega = 7.5265$
Response: $R = 0.0574$ $\rho = 2.9680$



Driven well above resonance


OscillIt
FDHO Web
Simulations

Initial Amplitude & Phase: $A = 0.3981$ $\alpha = -1.5708$
Stimulus: $A_s = 1.0000$ $\omega = 6.2832$
Response: $R = 0.3979$ $\rho = 1.5708$









Driven on resonance

Stimulus: $A_s = 1.0000$ $\omega = 5.0265$
Response: $R = 0.0697$ $\rho = 0.1405$



Driven well below resonance

Stimulus: $A_s = 1.0000$ $\omega = 7.5265$
 Response: $R = 0.0574$ $\rho = 2.9680$



Lorentz-Green's Function for high quality *FDHO*




Fig. 2.2.9 Beat formation.

Transient phasor $z_{\text{transient}}$ catches up with F -phasor and passes it.

OscillIt Web Simulation
 Beating ($\Gamma=0$)
 with transient and steady state

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions



Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Oscillator figures of merit: quality factors Q and $q=2\pi Q$

$$AAF = \frac{\text{Resonant response}}{\text{DC response}} = \frac{|G_{\omega_0}(\omega_s = \omega_0)|}{|G_{\omega_0}(0)|} = \frac{1/(2\Gamma\omega_0)}{1/\omega_0^2} = \frac{\omega_0}{2\Gamma} \equiv q \quad (\text{angular quality factor})$$

$$\text{Amplification factor } q = \omega_0/2\Gamma$$

Natural oscillation frequency is approximately $\nu_0 = \omega_0/2\pi$ (for $\omega_0 \gg \Gamma$ we have $\omega_0 \sim \omega_\Gamma$).

Oscillator figures of merit: quality factors Q and $q=2\pi Q$

$$AAF = \frac{\text{Resonant response}}{\text{DC response}} = \frac{|G_{\omega_0}(\omega_s = \omega_0)|}{|G_{\omega_0}(0)|} = \frac{1/(2\Gamma\omega_0)}{1/\omega_0^2} = \frac{\omega_0}{2\Gamma} \equiv q \quad (\text{angular quality factor})$$

$$\text{Amplification factor } q = \omega_0/2\Gamma$$

Natural oscillation frequency is approximately $\nu_0 = \omega_0/2\pi$ (for $\omega_0 \gg \Gamma$ we have $\omega_0 \sim \omega_\Gamma$).

$$\left(\begin{array}{l} t_{5\%} = 3/\Gamma = \text{Lifetime} \\ \text{for decaying oscillator} \\ \text{to lose 95\% of} \\ \text{amplitude} \end{array} \right) \text{times} \left(\nu_0 = \frac{\omega_0}{2\pi} \right) = \begin{array}{l} \text{number } n_{5\%} \\ \text{of oscillations} \\ \text{in a } t_{5\%} \text{ Lifetime} \end{array}$$

Oscillator figures of merit: quality factors Q and $q=2\pi Q$

$$AAF = \frac{\text{Resonant response}}{\text{DC response}} = \frac{|G_{\omega_0}(\omega_s = \omega_0)|}{|G_{\omega_0}(0)|} = \frac{1/(2\Gamma\omega_0)}{1/\omega_0^2} = \frac{\omega_0}{2\Gamma} \equiv q \quad (\text{angular quality factor})$$

$$\text{Amplification factor } q = \omega_0/2\Gamma$$

Natural oscillation frequency is approximately $\nu_0 = \omega_0/2\pi$ (for $\omega_0 \gg \Gamma$ we have $\omega_0 \sim \omega_\Gamma$).

$$\left(\begin{array}{l} t_{5\%} = 3/\Gamma = \text{Lifetime} \\ \text{for decaying oscillator} \\ \text{to lose 95\% of} \\ \text{amplitude} \end{array} \right) \text{times} \left(\nu_0 = \frac{\omega_0}{2\pi} \right) = \begin{array}{l} \text{number } n_{5\%} \\ \text{of oscillations} \\ \text{in a } t_{5\%} \text{ Lifetime} \end{array}$$

$$n_{5\%} = t_{5\%} \nu_0 = \frac{3}{\Gamma} \cdot \frac{\omega_0}{2\pi} \cong \frac{\omega_0}{2\Gamma} = q$$

The “Heartbeat Count”
measure of lifetime

Oscillator figures of merit: quality factors Q and $q=2\pi Q$

$$AAF = \frac{\text{Resonant response}}{\text{DC response}} = \frac{|G_{\omega_0}(\omega_s = \omega_0)|}{|G_{\omega_0}(0)|} = \frac{1/(2\Gamma\omega_0)}{1/\omega_0^2} = \frac{\omega_0}{2\Gamma} \equiv q \quad (\text{angular quality factor})$$

Amplification factor $q = \omega_0/2\Gamma$

Natural oscillation frequency is approximately $\nu_0 = \omega_0/2\pi$ (for $\omega_0 \gg \Gamma$ we have $\omega_0 \sim \omega_\Gamma$).

$$\left(\begin{array}{l} t_{5\%} = 3/\Gamma = \text{Lifetime} \\ \text{for decaying oscillator} \\ \text{to lose 95\% of} \\ \text{amplitude} \end{array} \right) \text{times} \left(\nu_0 = \frac{\omega_0}{2\pi} \right) = \begin{array}{l} \text{number } n_{5\%} \\ \text{of oscillations} \\ \text{in a } t_{5\%} \text{ Lifetime} \end{array}$$

$$n_{5\%} = t_{5\%} \nu_0 = \frac{3}{\Gamma} \cdot \frac{\omega_0}{2\pi} \cong \frac{\omega_0}{2\Gamma} = q$$

The “Heartbeat Count”
measure of lifetime

Energy decay
(proportional to the square of oscillator amplitude): $(e^{\Gamma t})^2 = e^{-2\Gamma t} \quad dE = -2\Gamma E$

Oscillator figures of merit: quality factors Q and $q=2\pi Q$

$$AAF = \frac{\text{Resonant response}}{\text{DC response}} = \frac{|G_{\omega_0}(\omega_s = \omega_0)|}{|G_{\omega_0}(0)|} = \frac{1/(2\Gamma\omega_0)}{1/\omega_0^2} = \frac{\omega_0}{2\Gamma} \equiv q \quad (\text{angular quality factor})$$

$$\text{Amplification factor } q = \omega_0/2\Gamma$$

Natural oscillation frequency is approximately $\nu_0 = \omega_0/2\pi$ (for $\omega_0 \gg \Gamma$ we have $\omega_0 \sim \omega_\Gamma$).

$$\left(\begin{array}{l} t_{5\%} = 3/\Gamma = \text{Lifetime} \\ \text{for decaying oscillator} \\ \text{to lose 95\% of} \\ \text{amplitude} \end{array} \right) \text{times} \left(\nu_0 = \frac{\omega_0}{2\pi} \right) = \begin{array}{l} \text{number } n_{5\%} \\ \text{of oscillations} \\ \text{in a } t_{5\%} \text{ Lifetime} \end{array}$$

$$n_{5\%} = t_{5\%} \nu_0 = \frac{3}{\Gamma} \cdot \frac{\omega_0}{2\pi} \cong \frac{\omega_0}{2\Gamma} = q$$

The “Heartbeat Count”
measure of lifetime

Energy decay

$$\text{(proportional to the square of oscillator amplitude): } \left(e^{\Gamma t} \right)^2 = e^{-2\Gamma t} \quad dE = -2\Gamma E$$

Relative amount

of energy lost

$$\text{each cycle period} = \tau_0 \left(\frac{-dE}{E} \right) = \frac{2\Gamma}{\nu_0} \equiv \frac{1}{Q} = \frac{2\pi}{q}$$

$$\left(\tau_0 = \frac{1}{\nu_0} \right)$$

$$Q = (\text{Standard angular quality factor}) = \frac{q}{2\pi}$$

Oscillator figures of merit: Uncertainty 1/q

To see a beat we need $\tau_{\text{half-beat}}$ to be less than $\tau_{5\%}$ or $3/\Gamma$. (Here we approximate $\pi \sim 3.0$, again.)

$$\pi / |\omega_s - \omega_0| < 3 / \Gamma$$

$$|\omega_s - \omega_0| > \Gamma$$

This means ω -detuning error is greater than or equal to the decay rate Γ .

Any detuning less than Γ is virtually undetectable.

Total ω uncertainty is $\pm\Gamma$ or twice Γ (that is: FWHM $\Delta\omega = 2\Gamma$). Linear frequency uncertainty is:

The *relative frequency uncertainty* $\frac{2\Gamma}{\omega_0} = \frac{\Delta\omega}{\omega_0} = \frac{1}{q} = \frac{\Delta\nu}{\nu_0}$ $\Delta\nu = \Delta\omega / 2\pi = \Gamma / \pi$

is the *inverse* of the *angular quality factor* q .

If we think of the 5% or 4.321% lifetime of a musical note as its time uncertainty Δt , then:

$$\Delta t \Delta \nu = 3 / \pi \approx 1$$

$$\Delta t = t_{5\%} = 3 / \Gamma$$

$$\Delta t = t_{4.321\%} = \pi / \Gamma$$

Very precise measures of imprecision

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

➔ *Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)*

Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Approximate Lorentz-Green's Function for high quality *FDHO* (Quantum propagator)

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} \xrightarrow{\omega_s \rightarrow \omega_0} \frac{1}{2\omega_s} \frac{1}{\omega_0 - \omega_s - i\Gamma} \approx \frac{1}{2\omega_0} \frac{1}{\Delta - i\Gamma} = \frac{1}{2\omega_0} L(\Delta - i\Gamma)$$

Complex detuning-decay $\delta = \Delta - i\Gamma$ variable δ is defined with the real detuning $\Delta = \omega_0 - \omega_s$

Approximate Lorentz-Green's Function for high quality *FDHO* (Quantum propagator)

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} \xrightarrow{\omega_s \rightarrow \omega_0} \frac{1}{2\omega_s} \frac{1}{\omega_0 - \omega_s - i\Gamma} \approx \frac{1}{2\omega_0} \frac{1}{\Delta - i\Gamma} = \frac{1}{2\omega_0} L(\Delta - i\Gamma)$$

Complex detuning-decay $\delta = \Delta - i\Gamma$ variable δ is defined with the real detuning $\Delta = \omega_0 - \omega_s$

$$L(\Delta - i\Gamma) = \frac{1}{\Delta - i\Gamma} = \text{Re } L + i \text{Im } L = \frac{\Delta}{\Delta^2 + \Gamma^2} + i \frac{\Gamma}{\Delta^2 + \Gamma^2} = |L|^2 \Delta + i |L|^2 \Gamma$$

Approximate Lorentz-Green's Function for high quality *FDHO* (Quantum propagator)

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} \xrightarrow{\omega_s \rightarrow \omega_0} \frac{1}{2\omega_s} \frac{1}{\omega_0 - \omega_s - i\Gamma} \approx \frac{1}{2\omega_0} \frac{1}{\Delta - i\Gamma} = \frac{1}{2\omega_0} L(\Delta - i\Gamma)$$

Complex detuning-decay $\delta = \Delta - i\Gamma$ variable δ is defined with the real detuning $\Delta = \omega_0 - \omega_s$

$$\begin{aligned} L(\Delta - i\Gamma) &= \frac{1}{\Delta - i\Gamma} = \text{Re } L + i \text{Im } L = \frac{\Delta}{\Delta^2 + \Gamma^2} + i \frac{\Gamma}{\Delta^2 + \Gamma^2} = |L|^2 \Delta + i |L|^2 \Gamma \\ &= |L| e^{i\rho} = |L| \cos \rho + i |L| \sin \rho = \frac{\cos \rho}{\sqrt{\Delta^2 + \Gamma^2}} + i \frac{\sin \rho}{\sqrt{\Delta^2 + \Gamma^2}} \text{ where: } |L| = \frac{1}{\sqrt{\Delta^2 + \Gamma^2}} \end{aligned}$$

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)

➔ Common Lorentzian (a.k.a. Witch of Agnesi)

Smith Charts

Approximate Lorentz-Green's Function for high quality *FDHO* (Quantum propagator)

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} \xrightarrow{\omega_s \rightarrow \omega_0} \frac{1}{2\omega_s} \frac{1}{\omega_0 - \omega_s - i\Gamma} \approx \frac{1}{2\omega_0} \frac{1}{\Delta - i\Gamma} = \frac{1}{2\omega_0} L(\Delta - i\Gamma)$$

Complex detuning-decay $\delta = \Delta - i\Gamma$ variable δ is defined with the real detuning $\Delta = \omega_0 - \omega_s$

$$L(\Delta - i\Gamma) = \frac{1}{\Delta - i\Gamma} = \text{Re } L + i \text{Im } L = \frac{\Delta}{\Delta^2 + \Gamma^2} + i \frac{\Gamma}{\Delta^2 + \Gamma^2} = |L|^2 \Delta + i |L|^2 \Gamma$$

$$= |L| e^{i\rho} = |L| \cos \rho + i |L| \sin \rho = \frac{\cos \rho}{\sqrt{\Delta^2 + \Gamma^2}} + i \frac{\sin \rho}{\sqrt{\Delta^2 + \Gamma^2}} \text{ where: } |L| = \frac{1}{\sqrt{\Delta^2 + \Gamma^2}}$$

Ideal Lorentz-Green's functions


$$L = \frac{\Delta + i\Gamma}{\Delta^2 + \Gamma^2} = |L| e^{i\rho}$$

Inverse decay rate $\frac{1}{\Gamma}$ axis
(Lifetime)

Smith plots

$$|L| = \frac{1}{\Gamma} \sin \rho$$

$$|L| = \frac{1}{\Delta} \cos \rho$$



Approximate Lorentz-Green's Function for high quality *FDHO* (Quantum propagator)

$$G_{\omega_0}(\omega_s) = \frac{1}{\omega_0^2 - \omega_s^2 - i2\Gamma\omega_s} \xrightarrow{\omega_s \rightarrow \omega_0} \frac{1}{2\omega_s} \frac{1}{\omega_0 - \omega_s - i\Gamma} \approx \frac{1}{2\omega_0} \frac{1}{\Delta - i\Gamma} = \frac{1}{2\omega_0} L(\Delta - i\Gamma)$$

Complex detuning-decay $\delta = \Delta - i\Gamma$ variable δ is defined with the real detuning $\Delta = \omega_0 - \omega_s$

$$L(\Delta - i\Gamma) = \frac{1}{\Delta - i\Gamma} = \text{Re } L + i \text{Im } L = \frac{\Delta}{\Delta^2 + \Gamma^2} + i \frac{\Gamma}{\Delta^2 + \Gamma^2} = |L|^2 \Delta + i |L|^2 \Gamma$$

$$= |L| e^{i\rho} = |L| \cos \rho + i |L| \sin \rho = \frac{\cos \rho}{\sqrt{\Delta^2 + \Gamma^2}} + i \frac{\sin \rho}{\sqrt{\Delta^2 + \Gamma^2}} \text{ where: } |L| = \frac{1}{\sqrt{\Delta^2 + \Gamma^2}}$$




Fig. 2.2.13 Ideal Lorentzian in inverse rate space. (Smith life-time $1/\Gamma$ vs. beat-period $1/\Delta$ coordinates)

Constant Δ and Γ curves in Fig. 2.2.13 are orthogonal circles of $1/z$ -dipolar coordinates. Recall Fig. 1.10.11.

1D forced-damped-harmonic oscillator equations and Green's function solutions

Linear harmonic oscillator equation of motion.

Linear damped-harmonic oscillator equation of motion.

Frequency retardation and amplitude damping

Figure of oscillator merit (the 5% solution $3/\Gamma$ and other numbers)

Linear forced-damped-harmonic oscillator equation of motion.

Properties of Green's function solutions and their mathematical/physical behavior

Phase lag and amplitude resonance amplification

Figure of resonance merit: Quality factor $q = \omega_0/2\Gamma$

Complete Green's Solution for the FDHO (Forced-Damped-Harmonic Oscillator)

Transient solutions vs. Steady State solutions

Quality factors: Beat, lifetimes, and uncertainty

Approximate Lorentz-Green's Function for high quality FDHO (Quantum propagator)


Common Lorentzian (a.k.a. Witch of Agnesi) ←

→ Smith Charts


An FDHO Green's
Function
Slide rule

A plot of
 $f(z) = 1/z$

For wavy
"Ohm's Laws"
 $V = I \cdot Z$
 $I = V/Z$



The Common Lorentzian (a.k.a. The Witch of Agnesi)



$$x^2 = b^2 \cot^2 \theta = b^2 \frac{\cos^2 \theta}{\sin^2 \theta} = b^2 \frac{1 - \sin^2 \theta}{\sin^2 \theta} = \frac{b^2}{\sin^2 \theta} - b^2$$

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{b}{y} \quad y = \frac{b}{x^2 + b^2}$$

Common Lorentzian function I.
(imaginary "absorbive" part)

Maria Gaetana Agnesi





Born May 16, 1718
Died January 9, 1799 (aged 80)
Residence Italy
Nationality Italy
Fields Mathematics

The Common Lorentzian (a.k.a. The Witch of Agnesi)

Maria Gaetana Agnesi



Born May 16, 1718
Died January 9, 1799 (aged 80)
Residence Italy
Nationality Italy
Fields Mathematics



$$x^2 = b^2 \cot^2 \theta = b^2 \frac{\cos^2 \theta}{\sin^2 \theta} = b^2 \frac{1 - \sin^2 \theta}{\sin^2 \theta} = \frac{b^2}{\sin^2 \theta} b^2$$


$$\frac{x}{y} = \frac{b \cot \theta}{(1/b) \cos \theta \sin \theta} = \frac{b^2 \cos \theta}{\cos \theta \sin^2 \theta} = \frac{b^2}{\sin^2 \theta}$$

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{b}{y} \quad y = \frac{b}{x^2 + b^2}$$

Common Lorentzian function I.
(imaginary "absorbtive" part)

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{x}{y} \quad y = \frac{x}{x^2 + b^2}$$

Common Lorentzian function II.
(real "refractory" part)





The Common Lorentzian (a.k.a. The Witch of Agnesi)

Maria Gaetana Agnesi



Born May 16, 1718
Died January 9, 1799 (aged 80)
Residence Italy
Nationality Italy
Fields Mathematics



$$x^2 = b^2 \cot^2 \theta = b^2 \frac{\cos^2 \theta}{\sin^2 \theta} = b^2 \frac{1 - \sin^2 \theta}{\sin^2 \theta} = \frac{b^2}{\sin^2 \theta} b^2$$


$$\frac{x}{y} = \frac{b \cot \theta}{(1/b) \cos \theta \sin \theta} = \frac{b^2 \cos \theta}{\cos \theta \sin^2 \theta} = \frac{b^2}{\sin^2 \theta}$$

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{b}{y} \quad y = \frac{b}{x^2 + b^2}$$

Common Lorentzian function I. (imaginary "absorbive" part)

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{x}{y} \quad y = \frac{x}{x^2 + b^2}$$

Common Lorentzian function II. (real "refractory" part)





The Common Lorentzian (a.k.a. The Witch of Agnesi)

Maria Gaetana Agnesi



Born May 16, 1718
 Died January 9, 1799 (aged 80)
 Residence Italy
 Nationality Italy
 Fields Mathematics




$$x^2 = b^2 \cot^2 \theta = b^2 \frac{\cos^2 \theta}{\sin^2 \theta} = b^2 \frac{1 - \sin^2 \theta}{\sin^2 \theta} = \frac{b^2}{\sin^2 \theta} - b^2$$

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{b}{y} \quad y = \frac{b}{x^2 + b^2}$$


Common Lorentzian function I.
(imaginary "absorbptive" part)

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{x}{y} \quad y = \frac{x}{x^2 + b^2}$$

Common Lorentzian function II.
(real "refractory" part)



Compare ideal Lorentzians ($\Gamma=0.2$) with a very non-ideal one ($\Gamma=2$)





The Common Lorentzian (a.k.a. The Witch of Agnesi)

Maria Gaetana Agnesi



Born May 16, 1718
 Died January 9, 1799 (aged 80)
 Residence Italy
 Nationality Italy
 Fields Mathematics




$$x^2 = b^2 \cot^2 \theta = b^2 \frac{\cos^2 \theta}{\sin^2 \theta} = b^2 \frac{1 - \sin^2 \theta}{\sin^2 \theta} = \frac{b^2}{\sin^2 \theta} b^2$$

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{b}{y} \quad \boxed{y = \frac{b}{x^2 + b^2}}$$

Common Lorentzian function I.
(imaginary "absorbive" part)

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{x}{y} \quad \boxed{y = \frac{x}{x^2 + b^2}}$$


Common Lorentzian function II.
(real "refractory" part)



Field:
 $f^*(z^*) = 1/z^{2*} = e^{i2\theta}/r^2$
 $\mathbf{F}(x,y) = (\cos 2\theta, \sin 2\theta)/r^2$
Potential:
 $\phi(z) = 1/z$
 $= (\cos \theta)/r + i(\sin \theta)/r$
 $= \Phi + i A$

Scalar potentials


$$\Phi = (a/r) \cos \theta = \text{const.}$$




Vector potentials

$$A = (a/r) \sin \theta = \text{const.}$$


Fig. 10.11 Dipole \mathbf{F} -field $f(z) = 1/z^2$ and scalar potential ($\Phi = \text{const.}$)-circles orthogonal to ($A = \text{const.}$)-circles.




From: Fig. 1.10.12



From: Fig. 1.10.12



From: Fig. 1.10.12




$$x^2 = b^2 \cot^2 \theta = b^2 \frac{\cos^2 \theta}{\sin^2 \theta} = b^2 \frac{1 - \sin^2 \theta}{\sin^2 \theta} = \frac{b^2}{\sin^2 \theta} b^2$$

$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{b}{y}$$

$$y = \frac{b}{x^2 + b^2}$$


*Common Lorentzian function I.
(imaginary "absorbitive" part)*



$$b = \frac{2}{5}$$

$$y = r \sin \theta$$

$$= (1/b) \cos \theta \sin \theta$$



$$x^2 + b^2 = \frac{b^2}{\sin^2 \theta} = \frac{x}{y}$$

$$y = \frac{x}{x^2 + b^2}$$
 Common Lorentzian function II.
 (real "refractory" part)

